

NUCLEAR DIMENSION AND n -COMPARISON

LEONEL ROBERT

ABSTRACT. It is shown that if a C^* -algebra has nuclear dimension n then its Cuntz semigroup has the property of n -comparison. It then follows from results by Ortega, Perera, and Rørdam that σ -unital C^* -algebras of finite nuclear dimension (and even of nuclear dimension at most ω) are stable if and only if they have no non-zero unital quotients and no non-zero bounded traces.

1. INTRODUCTION

In [7], Winter and Zacharias define nuclear dimension for C^* -algebras. This is a form of noncommutative dimension which directly generalizes the covering dimension of topological spaces. Finite nuclear dimension is specially relevant to the classification of C^* -algebras. The simple C^* -algebras of finite nuclear dimension have been proposed as a likely class of C^* -algebras for which Elliott's classification in terms of K-theory and traces holds true.

In the main result of this paper it is shown that the Cuntz semigroup of a C^* -algebra of finite nuclear dimension n satisfies the n -comparison property. For $n = 0$, this property is the same as almost unperforation in the Cuntz semigroup. For arbitrary n , it is reminiscent of the comparability between vector bundles whose fibrewise dimensions differ sufficiently relative to the dimension of the base space.

The n -comparison property for the Cuntz semigroup was first considered by Toms and Winter (see [6, Lemma 6.1]). They showed that n -comparison holds under the more restrictive assumption that the C^* -algebra is simple unital of decomposition rank n (the decomposition rank bounds the nuclear dimension, and, unlike nuclear dimension, is infinite for UCT Kirchberg algebras). The n -comparison property was subsequently studied, and more precisely defined, by Ortega, Perera and Rørdam in [5]. These authors obtained a simple criterion of stability for σ -unital C^* -algebras with n -comparison in their Cuntz semigroup: the C^* -algebra is stable if and only if it has no non-zero unital quotients and no non-zero bounded 2-quasitraces. By Theorem 1 below, this stability criterion applies to all C^* -algebras of finite nuclear dimension. It then follows that σ -unital C^* -algebras of finite nuclear dimension have the corona factorization property.

Let us recall the definition of nuclear dimension given in [7].

Definition 1. *The C^* -algebra A has nuclear dimension n if n is the smallest natural number for which there exist nets of completely positive contractions (henceforth abbreviated as c.p.c.)*

$$\psi_\lambda^i: A \rightarrow F_\lambda^i \text{ and } \phi_\lambda^i: F_\lambda^i \rightarrow A,$$

with $i = 0, 1, \dots, n$, $\lambda \in \Lambda$, and F_λ^i finite dimensional C^* -algebras for all i and λ , such that

- (i) ϕ_λ^i is an order 0 map (i.e., preserves orthogonality) for all i and λ ,
- (ii) $\lim_\lambda \sum_{i=1}^n \phi_\lambda^i \psi_\lambda^i(a) = a$ for all $a \in A$.

If no such n exists then A has infinite nuclear dimension.

Let us recall the definition given in [5] of the n -comparison property of an ordered semigroup. For x, y elements of an ordered semigroup S , let us write $x \leq_s y$ if $(k+1)x \leq ky$ for some $k \in \mathbb{N}$.

Definition 2. The ordered semigroup S has the n -comparison property if $x \leq_s y_i$ for $x, y_i \in S$ and $i = 0, 1, \dots, n$, implies $x \leq \sum_{i=0}^n y_i$.

Let $Cu(A)$ denote the stabilized Cuntz semigroup of the C^* -algebra A (i.e., the semigroup $W(A \otimes \mathcal{K})$; see [2]).

It is shown in Lemma 1 below that for $Cu(A)$ the n -comparison property can be reformulated as follows: if $[a], [b_i] \in Cu(A)$, with $i = 0, 1, \dots, n$, satisfy that for each i there is $\varepsilon_i > 0$ such that $d_\tau(a) \leq (1 - \varepsilon_i)d_\tau(b_i)$ for all dimension functions d_τ induced by lower semicontinuous 2-quasitraces, then $[a] \leq \sum_{i=0}^n [b_i]$. It is this formulation of the n -comparison property that is used by Toms and Winter in [6], and that may potentially have the most applications.

Theorem 1. If A has nuclear dimension n then $Cu(A)$ has the n -comparison property.

The following section is dedicated to the proof of Theorem 1. The last section discusses the application of Theorem 1, and of a variation on it that relates to ω -comparison, to establishing the stability of C^* -algebras of finite (or at most ω) nuclear dimension.

2. PROOF OF THEOREM 1

Let us start by proving that the property of n -comparison may be formulated using comparison by lower semicontinuous 2-quasitraces instead of the relation $\leq_s$. This result, however, will not be needed in the proof of Theorem 1.

For $[a], [b] \in Cu(A)$, elements of the Cuntz semigroup of A , let us write $[a] <_\tau [b]$ if there is $\varepsilon > 0$ such that $d_\tau(a) \leq (1 - \varepsilon)d_\tau(b)$ for all dimension functions d_τ induced by lower semicontinuous 2-quasitraces $\tau: A^+ \rightarrow [0, \infty]$ (see [3, Section 4]). We do not assume that the 2-quasitraces are necessarily finite on a dense subset of A^+ .

Lemma 1. The ordered semigroup $Cu(A)$ has the property of n -comparison if and only if for $[a], [b_i] \in Cu(A)$, with $i = 0, 1, \dots, n$, $[a] <_\tau [b_i]$ for all i implies that $[a] \leq \sum_{i=0}^n [b_i]$.

Proof. It is clear that if $[a] \leq_s [b]$ then $[a] <_\tau [b]$. Thus, n -comparison is implied by the property stated in the lemma. Suppose that we have n -comparison in $Cu(A)$. Let $[a], [b_i] \in Cu(A)$, with $i = 0, 1, \dots, n$, be such that $[a] <_\tau [b_i]$ for all i . Let us show that $[(a - \varepsilon)_+] \leq_s [b_i]$ for all $\varepsilon > 0$ and all i . Let $\lambda: Cu(A) \rightarrow [0, \infty]$ be additive and order preserving and let us define $\tilde{\lambda}: Cu(A) \rightarrow [0, \infty]$ by

$\tilde{\lambda}([c]) = \sup_{\delta > 0} \lambda([(c - \delta)_+])$. It is known that there is a lower semicontinuous 2-quasitrace τ such that $\tilde{\lambda}([a]) = d_\tau(a)$ for all $a \in (A \otimes \mathcal{K})^+$ (see [3, Proposition 4.2, Lemma 4.7]). We have

$$\lambda([(a - \varepsilon)_+]) \leq \tilde{\lambda}([a]) \leq (1 - \varepsilon_i) \tilde{\lambda}([b_i]) \leq (1 - \varepsilon_i) \lambda([b_i])$$

for all i . By [5, Proposition 2.1], this implies that $[(a - \varepsilon)_+] \leq_s [b_i]$ for all i . Since $Cu(A)$ has n -comparison, $[(a - \varepsilon)_+] \leq \sum_{i=0}^n [b_i]$. Taking supremum over $\varepsilon > 0$, we get that $[a] \leq \sum_{i=0}^n [b_i]$. $\square$

Throughout the rest of this section Λ denotes the index set in the definition of nuclear dimension. This set may be chosen to be the pairs (F, ε) where $F \subseteq A$ is finite and $\varepsilon > 0$. Let us denote by A_Λ the algebra $\prod_\lambda A / \bigoplus_\lambda A$. Let $\iota: A \rightarrow A_\Lambda$ denote the diagonal embedding of A into A_Λ .

Notation convention. For a family of C^* -algebras (A_λ) we use the notation $\bigoplus_\lambda A_\lambda$ to refer to the C^* -algebra of nets (x_λ) such that $\|x_\lambda\| \rightarrow 0$, while $\prod_\lambda A_\lambda$ denotes the nets of uniformly bounded norm.

In [7, Proposition 3.2] Winter and Zacharias show that if A has nuclear dimension n then the maps ψ_λ^i in the definition of nuclear dimension may be chosen asymptotically of order 0. That is, such that the induced maps $\psi^i: A \rightarrow \prod_\lambda F_\lambda^i / \bigoplus_\lambda F_\lambda^i$, for $i = 0, 1, \dots, n$, have order 0. We get the following proposition as a result of this.

Proposition 1. *If A has nuclear dimension n then for $i = 0, 1, \dots, n$ there are c.p.c. order 0 maps $\psi^i: A \rightarrow \prod_\lambda F_\lambda^i / \bigoplus_\lambda F_\lambda^i$ and $\phi^i: \prod_\lambda F_\lambda^i / \bigoplus_\lambda F_\lambda^i \rightarrow A_\Lambda$ such that*

$$(1) \quad \iota = \sum_{i=0}^n \phi^i \psi^i,$$

where $\iota: A \rightarrow A_\Lambda$ is the diagonal embedding of A into A_Λ .

Proof. As pointed out in the previous paragraph, by [7, Proposition 3.2] the maps ψ_λ^i in the definition of nuclear dimension may be chosen so that the induced maps ψ^i are of order 0.

The equation (1) is a consequence of Definition 1 (ii).

Let us show that the maps $\phi^i: \prod_\lambda F_\lambda^i / \bigoplus_\lambda F_\lambda^i \rightarrow A_\Lambda$, induced by the order 0 maps ϕ_λ^i , also have order 0. It is clear that $(\phi_\lambda^i): \prod_\lambda F_\lambda^i \rightarrow \prod_\lambda A$ has order 0. Hence, $\alpha \circ (\phi_\lambda^i): \prod_\lambda F_\lambda^i \rightarrow A_\Lambda$, where α is the quotient onto A_Λ , has order 0. We will be done once we show that if $\phi: C \rightarrow D$ is a c.p.c. map of order 0 and $\phi|_I = 0$ for some closed two sided ideal I , then the induced map $\tilde{\phi}: C/I \rightarrow D$ has order 0. By [8, Corollary 3.1], there is a $*$ -homomorphism $\pi: C \otimes C_0(0, 1] \rightarrow D$ such that $\phi(c) = \pi(c \otimes t)$ for all $c \in C$. From $\pi(I \otimes t) = 0$ we get that $\pi(I \otimes C_0(0, 1]) = 0$. Thus, π induces a $*$ -homomorphism $\tilde{\pi}: C/I \otimes C_0(0, 1] \rightarrow D$. Since $\tilde{\phi}(c) = \tilde{\pi}(c \otimes t)$ for all $c \in C/I$, $\tilde{\phi}$ has order 0. $\square$

An ordered semigroup S is said unperforated if $kx \leq ky$ for $x, y \in S$ and $k \in \mathbb{N}$ implies $x \leq y$.

Lemma 2. (i) If $Cu(A)$ is unperforated then so is $Cu(A/I)$ for any closed two-sided ideal I .

(ii) If $(A_\lambda)_\lambda$ are C^* -algebras such that $Cu(A_\lambda)$ is unperforated for all λ then so are $Cu(\prod_\lambda A_\lambda)$ and $Cu(\prod_\lambda A_\lambda / \bigoplus_\lambda A_\lambda)$.

Proof. (i) Let $[\tilde{a}], [\tilde{b}] \in Cu(A/I)$ be such that $k[\tilde{a}] \leq k[\tilde{b}]$ for some $k \in \mathbb{N}$. Then for $[a], [b] \in Cu(A)$ lifts of $[\tilde{a}]$ and $[\tilde{b}]$ we have

$$k[a] \leq k[b] + [i] \leq k([b] + [i]),$$

for some $i \in (I \otimes \mathcal{K})^+$ (by [1, Proposition 1]). Since $Cu(A)$ is unperforated, we have $[a] \leq [b] + [i]$, and passing to $Cu(A/I)$ we get that $[\tilde{a}] \leq [\tilde{b}]$.

(ii) Let $(a_\lambda)_\lambda, (b_\lambda)_\lambda \in (\prod_\lambda A_\lambda) \otimes \mathcal{K} \subseteq \prod_\lambda (A_\lambda \otimes \mathcal{K})$ be positive elements of norm at most 1 such that $k[(a_\lambda)_\lambda] \leq k[(b_\lambda)_\lambda]$ for some k . Let $\varepsilon > 0$. Then there is $\delta > 0$ such that

$$k[((a_\lambda - \varepsilon)_+)_\lambda] \leq k[((b_\lambda - \delta)_+)_\lambda].$$

Hence, $k[(a_\lambda - \varepsilon)_+] \leq k[(b_\lambda - \delta)_+]$ for all λ . Since $Cu(A_\lambda)$ is unperforated, $[(a_\lambda - \varepsilon)_+] \leq [(b_\lambda - \delta)_+]$. Let $x_\lambda \in A_\lambda \otimes \mathcal{K}$ be such that

$$(a_\lambda - 2\varepsilon)_+ = x_\lambda^* x_\lambda \text{ and } x_\lambda x_\lambda^* \in \text{her}((b_\lambda - \delta)_+).$$

Since $\|x_\lambda\|^2 \leq \|a_\lambda\| \leq 1$, we have $(x_\lambda)_\lambda \in \prod_\lambda (A_\lambda \otimes \mathcal{K})$. We also have $x_\lambda x_\lambda^* \leq \frac{1}{\delta} b_\lambda$, whence

$$((a_\lambda - 2\varepsilon)_+)_\lambda = (x_\lambda)_\lambda^* (x_\lambda)_\lambda \text{ and } (x_\lambda)_\lambda (x_\lambda)_\lambda^* \in \text{her}((b_\lambda)_\lambda).$$

But $(\prod_\lambda A_\lambda) \otimes \mathcal{K}$ sits as a hereditary subalgebra of $\prod_\lambda (A_\lambda \otimes \mathcal{K})$. Therefore, $(x_\lambda)_\lambda \in (\prod_\lambda A_\lambda) \otimes \mathcal{K}$ and so $(((a_\lambda - 2\varepsilon)_+)_\lambda) \leq [(b_\lambda)_\lambda]$. Since ε may be arbitrarily small, we have $[(a_\lambda)_\lambda] \leq [(b_\lambda)_\lambda]$ as desired. $\square$

Lemma 3. Let $a, b \in (A \otimes \mathcal{K})^+$. If $[\iota(a)] \leq [\iota(b)]$ then $[a] \leq [b]$.

Proof. Let $\varepsilon > 0$. Since $[\iota(a)] \leq [\iota(b)]$, there is $d \in A_\lambda \otimes \mathcal{K}$ such that $d^* \iota(b) d = \iota(a - \varepsilon)_+$. That is, there are $d_\lambda \in A_\lambda \otimes \mathcal{K}$ such that $d_\lambda^* b d_\lambda \rightarrow (a - \varepsilon)_+$. Thus, $[(a - \varepsilon)_+] \leq [b]$. Since $\varepsilon > 0$ may be arbitrarily small, we have $[a] \leq [b]$. $\square$

Remark 2. In proving Theorem 1, a stronger property than n -comparison will be shown to hold for C^* -algebras of nuclear dimension n : if $x, y_i \in Cu(A)$, with $i = 0, 1, \dots, n$, satisfy that $k_i x \leq k_i y_i$ for some $k_i \in \mathbb{N}$ and all i , then $x \leq \sum_{i=0}^n y_i$. This property, unlike n -comparison, does not seem to have a formulation in terms of comparison by lower semicontinuous 2-quasitraces.

Proof of Theorem 1. Suppose that there are $k_i \in \mathbb{N}$ such that $k_i[a] \leq k_i[b_i]$ for $i = 0, 1, \dots, n$. Since c.p.c. order 0 maps preserve Cuntz comparison (by [8, Corollary 3.5]), we have that $k_i[\psi^i(a)] \leq k_i[\psi^i(b_i)]$ for all i . Since the Cuntz semigroup of finite dimensional algebras is unperforated, we have by Lemma 2 that the Cuntz semigroup of $\prod_\lambda F_\lambda^i / \bigoplus_\lambda F_\lambda^i$ is unperforated. Thus, $[\psi^i(a)] \leq [\psi^i(b_i)]$. The maps ϕ^i preserve Cuntz equivalence (since they are c.p.c. of order 0), whence

$$[\phi^i \psi^i(a)] \leq [\phi^i \psi^i(b_i)] \leq [\sum_{j=0}^n \phi^j \psi^j(b_i)] = [\iota(b_i)].$$

So,

$$[\iota(a)] = \left[\sum_{i=0}^n \phi^i \psi^i(a) \right] \leq \sum_{i=0}^n [\phi^i \psi^i(a)] \leq \sum_{i=0}^n [\iota(b_i)].$$

By Lemma 3, this implies that $[a] \leq \sum_{i=0}^n [b_i]$. $\square$

3. STABILITY OF C^* -ALGEBRAS

A stable C^* -algebra has no non-zero unital quotients and no non-zero bounded 2-quasitraces (see [5, Proposition 4.6]). In [5, Proposition 4.8] Ortega, Perera and Rørdam show that the converse is true provided that the C^* -algebra is σ -unital and its Cuntz semigroup has the n -comparison property. This, combined with Theorem 1 and the fact that for exact C^* -algebras bounded 2-quasitraces are traces, implies that a σ -unital C^* -algebra of finite nuclear dimension is stable if and only if it has no non-zero unital quotients and no nonzero bounded traces. Ortega, Perera, and Rørdam also show that ω -comparison, a weakening of n -comparison, suffices to obtain the same stability criterion.

Definition 3. (c.f. [5, Definition 2.9]) *Let S be an ordered semigroup closed under the suprema of increasing sequences. Then S has the ω -comparison property if $x \leq_s y_i$, for $x, y_i \in S$ and $i = 0, 1, \dots$, implies $x \leq \sum_{i=0}^{\infty} y_i$.*

Remark 3. The definition of ω -comparison given above differs slightly from the definition given in [5]. Nevertheless, both definitions agree for ordered semigroups in the category Cu introduced in [2], and therefore, also for ordered semigroups arising as Cuntz semigroups of C^* -algebras.

A notion of nuclear dimension at most ω may be modelled after the statement of Proposition 1.

Definition 4. *Let us say that a C^* -algebra A has nuclear dimension at most ω if for $i = 0, 1, 2, \dots$ there are nets of c.p.c maps $\psi_{\lambda}^i: A \rightarrow F_{\lambda}^i$ and $\phi_{\lambda}^i: F_{\lambda}^i \rightarrow A$, with F_{λ}^i finite dimensional C^* -algebras, such that*

(ii) the induced maps $\psi^i: A \rightarrow \prod_{\lambda} F_{\lambda}^i / \bigoplus_{\lambda} F_{\lambda}^i$ and $\phi^i: \prod_{\lambda} F_{\lambda}^i / \bigoplus_{\lambda} F_{\lambda}^i \rightarrow A_{\Lambda}$ are c.p.c. order 0,

(iii) $\iota(a) = \sum_{i=0}^{\infty} \phi^i \psi^i(a)$ for all $a \in A$ (the sum on the right side is norm convergent).

For example, if the C^* -algebras $(A_i)_{i=0}^{\infty}$ all have finite nuclear dimension, then $\bigoplus_{i=0}^{\infty} A_i$ has nuclear dimension at most ω . It is not clear whether the assumption that the maps ψ_{λ}^i be asymptotically order 0 may be dropped in Definition 4 (and then proved), or if the other results on finite nuclear dimension proved in [7] also hold for nuclear dimension at most ω .

The proof of Theorem 1 goes through, mutatis mutandis, for nuclear dimension at most ω . We thus have,

Theorem 4. *If A has nuclear dimension at most ω then $Cu(A)$ has the ω -comparison property.*

Combined with the results of [5], Theorem 4 yields the following corollary, which improves on [4, Theorem 0.1], [5, Corollary 4.10] and [5, Corollary 5.13].

Corollary 1. *Let A be a C^* -algebra of nuclear dimension at most ω and let $B \subseteq A \otimes \mathcal{K}$ be hereditary and σ -unital. Then B is stable if and only if it has no non-zero unital quotients and no non-zero bounded traces. B has the corona factorization property.*

Proof. By Theorem 4, $Cu(A)$ has the ω -comparison property. Since $Cu(B)$ is an ordered semigroup ideal of $Cu(A)$, $Cu(B)$ has the ω -comparison property too. Hence, by [5, Proposition 4.8], B is stable if and only if it has no non-zero unital quotients and no non-zero bounded traces.

Let $C \subseteq B \otimes \mathcal{K}$ be full and hereditary, and suppose that $M_n(C)$ is stable. Then $M_n(C)$, and consequently C , cannot have non-zero unital quotients or bounded traces. Thus C is stable. This shows that B has the corona factorization property. $\square$

Remark 5. In the hypotheses of [5, Proposition 4.8] the C^* -algebra A is assumed separable. A closer look into the proof of this result reveals that it suffices to assume that the hereditary subalgebra $B \subseteq A \otimes \mathcal{K}$ be σ -unital. This justifies the application of [5, Proposition 4.8] in the above proof.

Acknowledgment. I am grateful to Wilhelm Winter for pointing out a correction to the proof of Lemma 2.

REFERENCES

- [1] A. Ciuperca, L. Robert, and L. Santiago, *The Cuntz semigroup of ideals and quotients and a generalized Kasparov Stabilization Theorem*, J. Operator Theory, to appear.
- [2] K. T. Coward, G. A. Elliott, and C. Ivanescu, *The Cuntz semigroup as an invariant for C^* -algebras*, J. Reine Angew. Math. **623** (2008), 161–193.
- [3] G. A. Elliott, L. Robert, and L. Santiago, *The cone of lower semicontinuous traces on a C^* -algebra* (2008), available at <http://arxiv.org/abs/0805.3122>.
- [4] P. W. Ng and W. Winter, *Nuclear dimension and the corona factorization property* (2009), available at <http://arxiv.org/abs/0904.0716>.
- [5] E. Ortega, F. Perera, and M. Rørdam, *The Corona Factorization property, Stability, and the Cuntz semigroup of a C^* -algebra* (2009), available at <http://arxiv.org/abs/0903.2917>.
- [6] A. S. Toms and W. Winter, *The Elliott conjecture for Villadsen algebras of the first type*, J. Funct. Anal. **256** (2009), no. 5, 1311–1340.
- [7] W. Winter and J. Zacharias, *The nuclear dimension of C^* -algebras* (2009), available at <http://arxiv.org/abs/0903.4914>.
- [8] ———, *Completely positive maps of order zero*, Münster J. of Math. **2** (2009), 311–324.

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, CANADA M3J 1P3

E-mail address: lrobert@mathstat.yorku.ca